

Electroweak Baryogenesis and Neutron Electric Dipole Moment from Supersymmetry ^{*}

Mayumi AOKI [†]

Graduate School of Humanities and Sciences

Ochanomizu University, Bunkyo-ku, Tokyo 112-8610, JAPAN

amayumi@fs.cc.ocha.ac.jp

Noriyuki OSHIMO

Institute for Cosmic Ray Research

University of Tokyo, Tanashi, Tokyo 188-0002, JAPAN

oshimo@icrr.u-tokyo.ac.jp

Akio SUGAMOTO

Department of Physics

Ochanomizu University, Bunkyo-ku, Tokyo 112-8610, JAPAN

sugamoto@phys.ocha.ac.jp

Abstract

Baryogenesis at the electroweak phase transition is discussed within the framework of the supersymmetric standard model. Implications of baryon asymmetry for the electric dipole moment of the neutron are also studied.

^{*}To be published in the Proceedings of the 3rd RESCEU International Symposium on Particle Cosmology.

[†]Research Fellow of the Japan Society for the Promotion of Science.

Astronomical observations indicate that there exist more baryons than anti-baryons in our universe. The ratio of baryon number to entropy is observed at present to be $\rho_B/s = (2 - 9) \times 10^{-11}$. This asymmetry may have been generated at the electroweak phase transition of the universe, since all the necessary conditions for baryogenesis seem to be satisfied within the framework of electroweak theories. The standard model (SM), however, cannot account for the asymmetry quantitatively. In this note, we report the possibility of baryogenesis by the charge transport mechanism in the supersymmetric standard model (SSM) based on $N=1$ supergravity.

The SSM has several complex parameters in addition to the Yukawa coupling constants for quarks. We can take for new physical complex parameters, without loss of generality, a Higgsino mass parameter m_H for the bilinear term of the Higgs superfields in the superpotential and a dimensionless coupling constant A for the trilinear terms of scalar fields which break supersymmetry softly. We express these parameters as $m_H = |m_H| \exp(i\theta)$ and $A = |A| \exp(i\alpha)$. The complex phases θ and α become new origins of CP violation.

The new sources of CP violation in the SSM give contributions to the electric dipole moment (EDM) of the neutron at the one-loop level through diagrams in which charginos, neutralinos or gluinos propagate together with squarks. The experimental upper bound of the neutron EDM is approximately $1 \times 10^{-25} \text{ ecm}$. 1) If the CP -violating phase θ is of order unity, the neutron EDM receives a contribution dominantly from the chargino diagram. From the experimental constraint, the masses of squarks are predicted to be larger than 1 TeV, while the chargino masses can be of order 100 GeV. In this case, charginos can mediate the charge transport mechanism. 2) If the phase θ is sufficiently smaller than unity, the chargino contribution to the EDM is negligible. The neutron EDM receives dominant contributions from the gluino and neutralino diagrams, leading to relaxed constraints on the masses of R -odd particles. The squarks are allowed to have masses of order 100 GeV, even if the phase α is of order unity. Then, t squarks assume the role of the mediators for the charge transport mechanism.

At the electroweak phase transition, if it is first order, bubbles of the broken phase nucleate in the symmetric phase. The charged gauginos λ and the charged Higgsinos ψ are in mass eigenstates in the symmetric phase. In the broken phase and the bubble wall, they are mixed to form charginos as mass eigenstates, owing to non-vanishing vacuum expectation values (vevs) v_1 and v_2 of Higgs bosons. At

the bubble wall, λ coming from the symmetric phase can be reflected to become ψ , and vice versa. The charginos in the broken phase can be transmitted to the symmetric phase and become λ or ψ . In these processes, CP violation is induced by the phase θ contained in the chargino mass matrix, making differences in reflection and transmission probabilities between CP -conjugate reactions. A net flux of hypercharge is then emitted into the symmetric phase. In the symmetric phase, the chemical potential of baryon number is related to a hypercharge density through equilibrium conditions. The induced hypercharge density in front of the wall makes the chemical potential of baryon number non-vanishing, which becomes a bias favoring a non-vanishing value of the baryon number. Baryogenesis then occurs, since the baryon number can change through an electroweak anomaly at a non-negligible rate, $\Gamma = 3\kappa(\alpha_W T)^4$, where T denotes a temperature at the phase transition and $\kappa \sim 0.1 - 1$. We take $\kappa = 1$ for numerical evaluations.

We calculated the baryon asymmetry for $\theta \sim 1$ and the chargino masses of order 100 GeV. The obtained ratio is $\rho_B/s \sim 10^{-10} - 10^{-11}$, which is compatible with the observed value. The ratio v_2/v_1 does not affect much the result. The velocity and width of the bubble wall are taken to be $0.1-1$ and $1/T - 5/T$, respectively. If the phase θ becomes of order 0.1, the resultant ratio is at most of order 10^{-11} . In the parameter ranges where a sufficient amount of the baryon asymmetry is produced, the neutron EDM is consistent with its experimental constraint, if the squark masses $M_{\tilde{q}}$ are larger than 1 TeV. For $1 \text{ TeV} \lesssim M_{\tilde{q}} \lesssim 10 \text{ TeV}$, the neutron EDM is predicted to be $10^{-25} - 10^{-26} \text{ ecm}$.

The baryon asymmetry can also be induced by t squarks. At the electroweak phase transition, the left-handed t squark $\tilde{t}_L$ and the right-handed t squark $\tilde{t}_R$ are in mass eigenstates in the symmetric phase, while non-vanishing vevs cause mixings of $\tilde{t}_L$ and $\tilde{t}_R$ in the bubble wall and the broken phase. At the bubble wall, $\tilde{t}_L$ coming from the symmetric phase can be reflected to become $\tilde{t}_R$, and vice versa. The mass eigenstates of t squarks in the broken phase can be transmitted to the symmetric phase and become $\tilde{t}_L$ or $\tilde{t}_R$. In these processes, the complex phase α in the mass squared-matrix of top squarks induces CP violation, causing differences in reflection and transmission rates between CP -conjugate reactions. A net hypercharge flux is emitted into the symmetric phase. Then, similarly to the baryogenesis by charginos, the baryon asymmetry is produced.

We calculated the baryon asymmetry for $\theta \ll 1$, $\alpha \sim 1$, and the squark masses of order 100 GeV. The obtained ratio can be as large as 5×10^{-11} in the same ranges

for the velocity and width of the wall as those for the chargino case. The ratio v_2/v_1 must vary in the wall for CP violation to occur. If the phase α becomes of order 0.1, it is difficult to produce a sufficient degree of asymmetry. The magnitude of the neutron EDM is below the experimental upper bound if $500 \text{ GeV} \lesssim \tilde{m}_2$, $\tilde{m}_2$ being the $SU(2)$ gaugino mass. For $500 \text{ GeV} \lesssim \tilde{m}_2 \lesssim 1 \text{ TeV}$, the neutron EDM is predicted to be $10^{-25} - 10^{-26} \text{ ecm}$.

We have studied the possibility of baryogenesis in the SSM, assuming that the electroweak phase transition is strongly first order. The charge transport mechanism mediated by charginos or t squarks can generate the ratio of baryon number to entropy consistent with its observed value, provided that the relevant CP -violating phase is not much suppressed and the relevant particles have masses of order 100 GeV. Then it is likely that the neutron EDM becomes $10^{-25} - 10^{-26} \text{ ecm}$. If the baryon asymmetry originates in the new source of CP violation in the SSM, the neutron EDM has a magnitude which can be explored in near-future experiments.

References

1. Aoki, M., Oshimo, N. and Sugamoto, A., 1997, *Prog. Theor. Phys.* **98**, 1179.
2. Aoki, M., Sugamoto, A. and Oshimo, N., 1997, *Prog. Theor. Phys.* **98**, 1325.